

Annapolis ENT Surgical Center, LLC

MEDICATIONS TO AVOID BEFORE SURGERY

ASPIRIN

Since aspirin can cause bleeding, it is important to avoid aspirin and medications which contain aspirin for at least 10 days before surgery and for one week after surgery.

Below is a list of common, brand-name prescription and over-the-counter drugs that contain aspirin:

Annapolis ENT Surgical Center, LLC

Aleve
APC Tablets
APC with Butalbital Tablets
APC with Codeine
Alka-Seltzer Effervescent
Alka-Seltzer Plus Cold Medicine
Anacin Analgesic Tablets
Anacin Analgesic Capsules
Arthritis Bayer Timed Release
Aspirin
Arthritis Strength Bufferin
Arthritis Strength BC Powder
Ascriptin with Codeine
Aspergum
Aspirin Suppositories
Aspirin with Codeine Tablets
Aspirin 325 mg. with Codeine
Axotal

Bayer Aspirin
Bayer Children's Chewable Tablets
Bayer Children's Cold Tablets
BC Powder
Buff-A Comp Tablets
Buff-A Comp No. 3 Tablets
Bufferin

Cama Arthritis Pain Reliever
Cama Inlay Tablets
Cetased, Improved
Congesprin Cold Tablets
Coricidin D Decongestant Tablets
Coricidin for Children
Coricidin Medilets Tablets for
Children
Coricidin-DC Capsules
Cosprin
Cosprin 650

Darvon with ASA
Darvon-N with ASA
Daypro
Dia-Gesic
Dihydrocodeine Compound Tablets
Dolprn #3
Dristan Decongestant Capsules and
Tablets

Easprin
8-Hour Bayer Timed-Release
Ecotrin Maximum Strength
Ecotrin Regular Strength
Ecotrin Tablets
Empirin
Empirin with Codeine
Emprazil Tablets
Emprazil-C Tablets EnTab
Enacaprin
Equagesic
Excedrin
Extra-Strength Ascriptin
Extra-Strength Bufferin

Extra-Strength Excedrin

4-Way Cold Tablets
Feldene
Fiorinal
Fiorinal with Codeine

Gemnisyn
Goody's Headache Powder

Hyco-Pap

Maximum Bayer Aspirin
Measurin Tablets
Mepro Compound Tablets
Methocarbamol with Aspirin Tablets
Midol
Momentum Muscular Backache
Formula

Norgesic & Norgesic Forte
Norwich Aspirin
Norwich Extra-Strength Aspirin

Oxaprozin
Oxycodone /Hydrocodone in
combination with Aspirin/Motrin
tablets

Pabirin Buffered Tablets
Panalgesic
Percodan & Percodan Demi Tablets
Piroxicam
Propoxyphene Compound 65

Quiet World Analgesic/Sleeping Aid

Robaxisal Tablets

SK-Oxycodone with Aspirin Tablets
SK-65 Compound Capsules
Sine-Off Sinus Medicine Tablets-
Aspirin Formula
Saletto
Soma Compound
Soma Compound with Codeine
St. Joseph Aspirin for Children
Supac
Synalgos Capsules
Synalgos-DC Capsules
Talwin Compound
Triaminicin Tablets

Ursinus Inlay-Tabs

Vanquish
Verin
Viro-Med Tablets

Zorprin

Annapolis ENT Surgical Center, LLC

Over-the-counter supplements, nutritional supplements and herbal and homeopathic medicines, especially **FISH OIL** and **VITAMIN E** must be stopped 10 days prior to surgery.

- The above is a list of common (but by no means all inclusive) aspirin containing medications.

IF IN DOUBT PLEASE CHECK WITH US!

- Please notify us immediately if you are currently taking anti-coagulant medication.
- Other non-aspirin medications which can cause bleeding and need to be stopped prior to surgery are:
 - Ibuprofen - stop 3-5 days prior to surgery
 - Motrin - stop 3-5 days prior to surgery
 - Advil - stop 3-5 days prior to surgery
 - Nuprin - stop 3-5 days prior to surgery
- **TYLENOL IS OKAY**

I HAVE READ AND UNDERSTAND THE ABOVE INFORMATION: _____

Patient's Signature

Date